

## Lattice and Semilattice Properties

Given a  $T_{3\frac{1}{2}}$  space  $(X, \tau)$ ,  $\mathcal{COM}(X)$  denotes the set described in [6], which consists of representatives of each  $T_2$  compactification class of  $X$ . In R2.6 and R2.7 of [6], Lubben's results that  $\mathcal{COM}(X)$  is a complete upper semi-lattice and that, if  $X$  is also locally compact,  $\mathcal{COM}(X)$  is a complete lattice are presented. As in [5],  $\mathcal{TB}(X)$  denotes the set of totally bounded uniformities for  $X$  which generate  $\tau$ . The map  $\Psi_0$  of [5], which associates  $\mathcal{U} \in \mathcal{TB}(X)$  with an equivalence class of compactifications, can easily be converted into an order isomorphism as follows.

**Definition R8.1**  $\Psi : \mathcal{TB}(X) \rightarrow \mathcal{COM}(X)$  is defined by letting  $\Psi(\mathcal{U})$  be the element of  $\mathcal{COM}(X)$  which represents the equivalence class  $\Psi_0(\mathcal{U})$ .

**Proposition R8.2**  $\mathcal{TB}(X)$  and  $\mathcal{COM}(X)$  are order isomorphic, with  $\Psi$  as an order isomorphism.

Proof: By R1.5 [5], for  $\mathcal{U}_1, \mathcal{U}_2$  in  $\mathcal{TB}(X)$ ,  $\mathcal{U}_1 \supseteq \mathcal{U}_2$  if and only if  $\Psi(\mathcal{U}_1) \geq \Psi(\mathcal{U}_2)$ . Thus  $\Psi$  is one-to-one and order preserving. By R1.4  $\Psi$  is onto.

Note that, by using R7.2.18 of [8] in a similar way, one could establish an order isomorphism between  $\mathcal{COM}(X)$  and the set of  $\mathcal{C}(\mathcal{U})$  for  $\mathcal{U}$  in  $\mathcal{TB}(X)$ , where  $\mathcal{C}(\mathcal{U})$  is the set of all real-valued, uniformly continuous maps from  $(X, \mathcal{U})$ .

In the following example  $\mathcal{COM}(X)$  is a complete lattice by R2.7, but  $\mathcal{TB}(X)$  (and so also  $\mathcal{COM}(X)$  by R8.2) is not even finitely distributive. Because in a lattice, either distributive law implies the other, the example only needs to verify that one fails.

**Example R8.3** Let  $(\mathbf{N}, \tau)$  be the positive intergers with the discrete topology, let  $E_4$  be equivalence mod 4, and let  $[x]_4$  denote the equivalence class of  $x$  mod 4. Let  $E_3$  be the equivalence relation generated by the partition  $\{[1]_4 \cup [4]_4, [2]_4 \cup [3]_4\}$ , let  $E_2$  be the equivalence relation generated by  $\{[1]_4 \cup [2]_4, [3]_4 \cup [4]_4\}$ , and let  $E_1$  be equivalence mod 2. Let  $\mathcal{V}_i = \mathcal{U}_m \vee \mathcal{U}_{E_i}$ , where  $\mathcal{U}_m$  and  $\mathcal{U}_{E_i}$  are the uniformities defined in subsection 2 of [7]. By R5.2.3,  $\mathcal{V}_i$  is in  $\mathcal{TB}(\mathbf{N})$  for  $i = 1, \dots, 4$ . For  $i = 1, 2, 3$  clearly  $E_4 \subseteq E_i$  and so  $\mathcal{V}_i \subseteq \mathcal{V}_4$ . Also  $E_2 \cap E_3 = E_4$  so that  $E_4 \in \mathcal{V}_2 \vee \mathcal{V}_3$ . As a result  $\mathcal{V}_2 \vee \mathcal{V}_3 = \mathcal{V}_4$  and  $\mathcal{V}_1 \wedge (\mathcal{V}_2 \vee \mathcal{V}_3) = \mathcal{V}_1 \neq \mathcal{U}_m$ . The next claim is that  $\mathcal{V}_1 \wedge \mathcal{V}_2 = \mathcal{V}_1 \wedge \mathcal{V}_3 = \mathcal{U}_m$ , which would show that  $\wedge$  is not distributive over  $\vee$ . Clearly  $\mathcal{U}_m \subseteq \mathcal{V}_1 \wedge \mathcal{V}_2$ . For the converse let  $V$  be in  $\mathcal{V}_1 \wedge \mathcal{V}_2$  and pick  $W \in \mathcal{V}_1 \wedge \mathcal{V}_2$  with  $W = W^{-1}$  and  $W \circ W \subseteq V$ . There is  $U \in \mathcal{U}_m$  such that  $U \cap E_1 \subseteq W$  and  $U \cap E_2 \subseteq W$ , and there is  $S \subseteq \mathbf{N}$  such that  $\mathbf{N} - S$  is finite and  $S \times S \subseteq U$ . Since  $V$  contains the diagonal,  $S \times S \subseteq V$  would imply  $V \in \mathcal{U}_m$ . Let  $(x, y) \in S \times S$ . If  $(x, y) \in E_1$  or  $(x, y) \in E_2$ , then  $(x, y) \in W \subseteq V$ . In the remaining case, pick  $t \in S$  such that  $(x, t) \in E_1$  but  $(x, t) \notin E_4$ . Then  $(x, t) \in U \cap E_1 \subseteq W$  and  $(t, y) \in U \cap E_2 \subseteq W$  and so  $(x, y) \in W \circ W \subseteq V$ . Similarly,  $\mathcal{V}_1 \wedge \mathcal{V}_3 = \mathcal{U}_m$ .

Chandler [1] presents a number of results focusing on this question: When is  $\mathcal{COM}(X)$  a complete lattice or a lattice? The next two lemmas lead to a sharpened version of R2.7. For both lemmas the following will be assumed:  $(Y, f)$  will denote a  $T_2$  compactification of  $X$  with  $y_1 \neq y_2$  in  $Y - f[X]$ . Let  $\mathcal{U} \in \mathcal{TB}(X)$  with  $\Psi_0(\mathcal{U}) = [(Y, f)]$ , and, making use of R7.1.1, let  $\mathcal{C} = \{g \in \mathcal{C}(\mathcal{U}) : G(y_1) = G(y_2) \text{ where } G \text{ is the unique continuous extension of } g \text{ to } Y\}$ . Finally let  $\mathcal{U}_1$  denote  $\mathcal{U}(\mathcal{C})$ , which by R7.2.7 and [4] is the smallest uniformity for  $X$  making every element of  $\mathcal{C}$  uniformly continuous.

**Lemma R8.4**  $\mathcal{U}_1 \in \mathcal{TB}(X)$ .

Proof: By R7.2.10 it is sufficient to show that  $\mathcal{C}$  separates points and closed sets of  $X$  in the sense of [8]. Let  $F$  be closed in  $X$  and let  $x \in X - F$ . Since  $f$  is an embedding,  $f(x) \notin c_Y(f[F])$ . Let  $S = c_Y(f[F]) \cup \{y_1, y_2\}$ , and pick  $W \in \mathcal{W}$ , the unique uniformity for  $Y$ , such that  $W[f(x)] \subseteq Y - S$ . There is a pseudo-metric  $d$  for  $Y$  such that  $W \in \mathcal{U}_d \subseteq \mathcal{W}$ . Then  $G(y) = \inf\{d(y, t) : t \in S\}$  is uniformly continuous by R7.2.14 and  $G[S] = \{0\}$  so that  $g = G \circ f$  is in  $\mathcal{C}$  and  $g[F] \subseteq \{0\}$ . Since  $W[f(x)]$  is a  $d$ -neighborhood of  $f(x)$ ,  $g(x) = G(f(x)) \neq 0$  and so  $\{g(x)\} \cap \overline{g[F]} = \emptyset$ , i.e.,  $\mathcal{C}$  separates points and closed sets of  $X$ .

**Lemma R8.5**  $\mathcal{U}_1$  is a proper subset of  $\mathcal{U}$ .

Proof: Since every element of  $\mathcal{C}$  is uniformly continuous from  $(X, \mathcal{U})$ , the smallest such uniformity,  $\mathcal{U}_1$ , is clearly contained in  $\mathcal{U}$ . Let  $\mathcal{W}$  be the unique uniformity for  $Y$ , let  $W_1 \in \mathcal{W}$  with  $W_1[y_1] \cap W_1[y_2] = \emptyset$ , and pick  $W \in \mathcal{W}$  with  $W = W^{-1}$  and  $W \circ W \circ W \subseteq W_1$ . Let  $(f \times f)^{-1}[W \cap f[X] \times f[X]] = U^*$ . Clearly  $U^* \in \mathcal{U}$  and it is sufficient to show  $U^* \notin \mathcal{U}_1$ . Deny the latter. Then there exist  $g_1, \dots, g_n$  in  $\mathcal{C}$  and  $\epsilon > 0$  such that

$$\bigcap_{i=1}^n (g_i \times g_i)^{-1}[V_\epsilon] \subseteq U^*.$$

Let  $\{a_t\}$  and  $\{b_r\}$  be nets in  $X$  such that  $f(a_t) \rightarrow y_1$  and  $f(b_r) \rightarrow y_2$ . With  $G_i$  denoting the continuous extension of  $g_i$  to  $Y$ , we have  $g_i(a_t) = G_i(f(a_t)) \rightarrow G_i(y_1) = G_i(y_2)$  and also  $g_i(b_r) = G_i(f(b_r)) \rightarrow G_i(y_2)$ . Using the definition of net convergence and the fact that  $W[y_i]$  is a neighborhood of  $y_i$ , we can find  $t$  and  $r$  such that  $f(a_t) \in W[y_1]$ ,  $f(b_r) \in W[y_2]$ , and  $|g_i(a_t) - g_i(b_r)| < \epsilon$  for all  $i$ . Then  $(a_t, b_r) \in U^*$  and so  $(f(a_t), f(b_r)) \in W$ . But then  $(y_1, y_2) \in W \circ W \circ W \subseteq W_1$ , a contradiction.

**Theorem R8.6** [Lubben]  $\mathcal{COM}(X)$  is a complete lattice if and only if  $X$  is locally compact.

Proof: The ‘if’ half of this result is R2.7. For the converse, it is sufficient to show that  $\mathcal{COM}(X)$  does not have a smallest element if  $X$  is not locally compact. What follows is a uniformity-oriented modification of Chandler’s argument for Theorem 2.19 in [1]. Let  $(Y, f)$  be a  $T_2$  compactification of  $X$  and let  $\mathcal{U} \in \mathcal{TB}(X)$  with  $\Psi_0(\mathcal{U}) = [(Y, f)]$ . Since  $X$  is not locally compact,  $f[X]$  is not open in  $Y$  and so there exist distinct points  $y_1, y_2$  in  $Y - f[X]$ . Let  $\mathcal{U}_1$  be defined as in the preceding lemmas. By R8.5 and R1.5  $\Psi(\mathcal{U}_1)$  is strictly smaller than  $\Psi(\mathcal{U})$ , the element of  $\mathcal{COM}(X)$  representing  $(Y, f)$ .

The next two results provide a condition under which a non-empty family in  $\mathcal{COM}(X)$  has an infimum. To do this, it is enough to focus on the isomorphic upper semi-lattice  $\mathcal{TB}(X)$ . Recall that any non-empty family of uniformities has an infimum, whose topology may be strictly smaller than the infimum (i.e., intersection) of the corresponding family of topologies generated by the uniformities.

**Lemma R8.7** Let  $\{\mathcal{U}_\alpha : \alpha \in \Delta\}$  be a non-empty family in  $\mathcal{TB}(X)$ .

Let  $\mathcal{U} = \wedge\{\mathcal{U}_\alpha : \alpha \in \Delta\}$ . Then  $\mathcal{U}$  is a totally bounded uniformity for  $X$  with  $\tau(\mathcal{U}) \subseteq \tau$  and  $\mathcal{C}(\mathcal{U}) \subseteq \cap\{\mathcal{C}(\mathcal{U}_\alpha) : \alpha \in \Delta\}$ .

Proof: This is clear from general facts about uniformities.

If one transfers the next result to  $\mathcal{COM}(X)$  via the order isomorphism  $\Psi$ , it provides a condition for the existence of an infimum of a family. This condition is a uniformity-oriented version of Chandler’s Theorem 2.18 in [1].

**Proposition R8.8** Let  $\{\mathcal{U}_\alpha : \alpha \in \Delta\}$  be a non-empty family in  $\mathcal{TB}(X)$ .

Let  $\mathcal{U} = \wedge\{\mathcal{U}_\alpha : \alpha \in \Delta\}$ . Then  $\mathcal{U} \in \mathcal{TB}(X)$  if and only if  $\cap\{\mathcal{C}(\mathcal{U}_\alpha) : \alpha \in \Delta\}$  separates points and closed sets.

Proof: First, let  $\mathcal{C} = \cap\{\mathcal{C}(\mathcal{U}_\alpha) : \alpha \in \Delta\}$  separate points and closed sets. By R7.2.10  $\mathcal{U}(\mathcal{C}) \in \mathcal{TB}(X)$  and, since  $\mathcal{C} \subseteq \mathcal{C}(\mathcal{U}_\alpha)$ , by R7.2.16  $\mathcal{U}(\mathcal{C}) \subseteq \mathcal{U}_\alpha$  for all  $\alpha$ . Thus  $\mathcal{TB}(X)$  contains a lower bound for  $\{\mathcal{U}_\alpha : \alpha \in \Delta\}$  and so  $\mathcal{U} \in \mathcal{TB}(X)$ . For the converse, the hypothesis implies by R7.2.15 that  $\mathcal{C}(\mathcal{U})$  separates points and closed sets, and clearly a larger family such as  $\mathcal{C}$  does also.

The next result shows that, if  $X$  is the rationals with the usual topology, then  $\mathcal{COM}(X)$  is not a lattice.

**Theorem R8.9** [Shirota 2] If  $X$  is first countable and not locally compact, then  $\mathcal{COM}(X)$  is not a lattice.

Proof: See Chandler [1], Theorem 5.12.

Finally,  $\mathcal{COM}(X)$  can have any infinite cardinality. Chandler [1] uses ideas from the proof of the following theorem to construct an example of a non-locally compact  $X$  for which  $\mathcal{COM}(X)$  is a lattice.

**Theorem R8.10** [Visliseni and Flaksmaier 3] Let  $\aleph$  be an infinite cardinal. Then there is a  $T_{3\frac{1}{2}}$  space  $(X, \tau)$  such that  $|\mathcal{COM}(X)| = \aleph$ .

Proof: See Chandler [1], Theorem 5.4.

Albert J. Klein 2004

<http://www.susanjkleinart.com/compactification/>

## References

An asterisk indicates a reference not seen by me.

1. Chandler, R. E., Hausdorff Compactifications, Marcel Dekker Inc., 1976
- 2\* Shirota, T., On Systems of Structures of a Completely Regular Space, Osaka Math. J. 41(1950), 131-143.
- 3\* Visliseni, Ju. and Flaksmaier, Ju., Power and Construction of the Structure of all Compact Extensions of a Completely Regular Space, Soviet Math. Dokl. 6(1965), 1423-1425.
4. Wilansky, A., Topology for Analysis, Ginn and Co., 1970.
5. This website, R1: Existence via Uniform Space Theory
6. This website, R2: Existence via Quotients of the Stone-Cěch Compactification
7. This website, R5: Finite-Point Compactifications
8. This website, R7: Uniform Continuity and Extension of Maps

## Added 2018

The following proposition is implicit in the proof of R8.6 given above.

**Proposition R8.Add.1** Let  $(X, \tau)$  be a  $T_{3\frac{1}{2}}$  space, which is not locally compact. Then the complete upper semi-lattice  $\mathcal{COM}(X)$  does not contain any minimal elements.

Proof: The argument for R8.6 shows that, given a  $T_2$  compactification of  $(X, \tau)$ , there is another which is a strictly smaller.

**Corollary R8.Add.2** Let  $(X, \tau)$  be a  $T_{3\frac{1}{2}}$  space, which is not locally compact. Then the complete upper semi-lattice  $\mathcal{TB}(X)$  does not contain any minimal elements.

Proof: This is immediate from R8.2 and the preceding proposition.

For example R8.Add.6 below, the fact that every uniformity contains a totally bounded uniformity generating the same topology is needed. Since that may not be widely known, an indication of its proof is provided in the next three items. The proof is adapted from the proximity approach to uniform ideas. An overview of proximity spaces and a bibliography of original sources are included in Nachman [10].

**Definition R8.Add.3** Let  $(X, \mathcal{U})$  be a uniform space. A proximal cover of  $X$  is a finite collection  $\{A_1, \dots, A_n\}$  of subsets of  $X$  for which there exist sets  $B_1, \dots, B_n$  and  $U \in \mathcal{U}$  such that  $X = \cup_{i=1}^n B_i$  and  $U[B_i] \subseteq A_i$  for  $1 \leq i \leq n$ .

Clearly, a proximal cover is a cover.

**Lemma R8.Add.4** Let  $(X, \mathcal{U})$  be a uniform space and let  $\{A_1, A_2\}$  be a proximal cover of  $X$ . Then there is a proximal cover  $\{D_1, D_2, D_3\}$  such that, with  $S = \cup_{i=1}^3 D_i \times D_i$ ,  $S \circ S \subseteq (A_1 \times A_1) \cup (A_2 \times A_2)$ .

Sketch of proof: Select  $B_1, B_2, U = U^{-1}$  for the proximal cover  $\{A_1, A_2\}$ . It follows easily that  $U[X - A_1] \subseteq X - B_1 \subseteq A_2$ . Pick  $V = V^{-1}$  in  $\mathcal{U}$  with  $V^6 \subseteq U$ . (Here the exponent indicates repeated composition, i.e.,  $V^2 = V \circ V$  etc.) Let  $D_1 = V^3[X - A_1]$ ,  $D_2 = V^6[X - A_1] - V[X - A_1]$ , and  $D_3 = X - V^4[X - A_1]$ . Note that  $D_1 \subseteq A_2$ ,  $D_2 \subseteq A_1 \cap A_2$ ,  $D_3 \subseteq A_1$ , and  $D_1 \cap D_3 = \emptyset$ . Let  $C_1 = V^2[X - A_1]$ ,  $C_2 = V^5[X - A_1] - V^2[X - A_1]$ , and  $C_3 = X - V^5[X - A_1]$ . Then  $X = \cup_{i=1}^3 C_i$  and  $V[C_i] \subseteq D_i$  for  $i \in \{1, 2, 3\}$  and so  $\{D_1, D_2, D_3\}$  is a proximal cover of  $X$ . With this choice of  $D_1, D_2, D_3$  the conclusion holds.

**Lemma R8.Add.5** Let  $(X, \mathcal{U})$  be a uniform space and let  $\mathcal{V}$  be defined as the set  $\{V \subseteq X \times X : \text{there is a proximal cover } \{A_1, \dots, A_n\} \text{ with } \cup_{i=1}^n A_i \times A_i \subseteq V\}$ . Then

- i)  $\mathcal{V}$  is a uniformity for  $X$ .
- ii)  $\mathcal{V}$  is totally bounded.
- iii)  $\mathcal{V} \subseteq \mathcal{U}$ .
- iv)  $\tau(\mathcal{U}) = \tau(\mathcal{V})$ .

Sketch of proof: For i): The diagonal, superset, and symmetry requirements (P2.1i,ii, and iv) are clear. Given proximal covers  $\{A_1, \dots, A_n\}$  and  $\{B_1, \dots, B_m\}$ , they determine another proximal cover  $\{A_i \cap B_j : 1 \leq i \leq n, 1 \leq j \leq m\}$ . The finite intersection requirement (P2.1iii) follows easily. For a proximal cover  $\{A_1, \dots, A_n\}$  and  $I \subseteq \{1, \dots, n\}$ , let  $A_I = \cup\{A_i : i \in I\}$  and  $B_I = \cup\{A_i : i \notin I\}$ . Then  $\{A_I, B_I\}$  is a proximal cover. Let  $V(I) = (A_I \times A_I) \cup (B_I \times B_I)$ . It is easy to check that  $\cup_{i=1}^n A_i \times A_i$  equals the finite intersection  $\cap\{V(I) : I \subseteq \{1, \dots, n\}\}$ . The triangle inequality (P2.1v) follows by applying R8.Add.4. For ii), let a proximal cover  $\{A_1, \dots, A_n\}$  be given and pick  $x_i \in A_i$ . Let  $V = \cup_{i=1}^n A_i \times A_i$ . Then  $\cup_{i=1}^n V[x_i] = \cup_{i=1}^n A_i = X$ . For iii) let a proximal cover  $\{A_1, \dots, A_n\}$  be given and let  $S = \cup_{i=1}^n A_i \times A_i$ . Let  $\{B_1, \dots, B_n\}$  and  $U \in \mathcal{U}$  be such that  $\cup_{i=1}^n B_i = X$  and  $U[B_i] \subseteq A_i$  for  $1 \leq i \leq n$ . Pick  $V = V^{-1}$  in  $\mathcal{U}$  with  $V \circ V \subseteq U$ . Let  $(x, y) \in V$ . For  $k$  with  $x \in B_k$ ,  $\{x, y\} \subseteq V[x] \subseteq V[V[B_k]] \subseteq U[B_k] \subseteq A_k$ . Thus  $V \subseteq S$ . For iv) note that iii) implies  $\tau(\mathcal{V}) \subseteq \tau(\mathcal{U})$ . Let  $x \in O \in \tau(\mathcal{U})$ . Pick  $U, V \in \mathcal{U}$  with  $U[x] \subseteq O$ ,  $V \circ V \subseteq U$ , and  $V = V^{-1}$ . Let  $A_1 = O$  and  $A_2 = V[X - V[x]]$ . The sets  $B_1 = V[x], B_2 = X - V[x]$  show that  $\{A_1, A_2\}$  is a proximal cover and so

$W = (A_1 \times A_1) \cup (A_2 \times A_2)$  is in  $\mathcal{V}$ . Then  $x \notin A_2$  so that  $W[x] = O$ . Thus  $O \in \tau(\mathcal{V})$ .

**Example R8.Add.6** Let  $\mathbf{R}$  be the real numbers with the usual topology and let  $\mathcal{U}_m$  be the uniformity for  $\mathbf{R}$  corresponding to the class of the one-point compactification. Let  $\mathbf{Q}$  be the set of rational numbers with the relative topology from  $\mathbf{R}$  and let  $\mathcal{U}$  be the subspace uniformity from  $\mathcal{U}_m$ . Then  $\mathcal{U}$  is a separated, totally bounded uniformity for  $\mathbf{Q}$  and so by R8.Add.2 there is  $\mathcal{V}$ , also a separated, totally bounded uniformity for  $\mathbf{Q}$ , such that  $\mathcal{V}$  is a proper subset of  $\mathcal{U}$ . Note that there is no uniformity  $\mathcal{W}$  for  $\mathbf{R}$  such that the subspace uniformity on  $\mathbf{Q}$  from  $\mathcal{W}$  is  $\mathcal{V}$ : Otherwise, by R8.Add.5 one could pick a totally bounded  $\mathcal{W}_1 \subseteq \mathcal{W}$ . Since  $\mathcal{U}_m \subseteq \mathcal{W}_1$ ,  $\mathcal{U}$  is contained in the subspace uniformity on  $\mathbf{Q}$  from  $\mathcal{W}$ . But  $\mathcal{V}$  is a proper subset of  $\mathcal{U}$ .

Comment: The construction in the previous example could be continued inductively to produce a decreasing sequence of separated totally bounded uniformities for  $\mathbf{Q}$  with the relative topology from  $\mathbf{R}$ . None of the uniformities in the sequence is the subspace uniformity of some uniformity for  $\mathbf{R}$ . Each determines a  $T_2$  compactification and so a uniform completion of  $\mathbf{Q}$ . It is not clear how interesting these might be.

### Additional References

9. Bentley, H.L., Herrlich, H., and Huček, M., The Historical Development of Uniform, Proximal, and Nearness Concepts in Topology, in: Handbook of the History of General Topology, vol. 2, C. E. Aull and R. Lowen, editors, Kulwer Academic Publishers, 1998.

10. Nachman, Louis Jack, Weak and Strong Constructions in Proximity Spaces, Thesis, The Ohio State University, Columbus, Ohio, 1968.

11. This website, P2: Uniform Spaces